

Infinite dimensional symmetry of corner transfer matrices

Brian Davies

SMS-118-93

CMA-MR55-93

Department of Mathematics,
School of Mathematical Sciences,
Australian National University,
Canberra, ACT 0200, Australia

December 1993

1991 *Mathematics subject classification*: primary 82B13; secondary 82B20.

Abstract. We review some of the recent developments in two dimensional statistical mechanics in which corner transfer matrices provide the vital link between the physical system and the representation theory of quantum affine algebras. This opens many new possibilities, because the eigenstates may be described using the properties of q -vertex operators.

1. Introduction

Transfer matrices have been a central object for the study of two-dimensional statistical mechanics systems for more than a half century. Originally the row transfer matrix was defined: it encapsulates the statistical information (via the Boltzmann weights) pertaining to the contribution from a single row of a regular lattice. Row transfer matrices are extremely complicated; for an Ising or six-vertex model on a square lattice of width L the row transfer matrix has dimension 2^L . Onsager's solution of the Ising model [1] used representation theory of finite-dimensional Lie algebras to transform the generators of the row transfer matrix to a tractable form. The next several major advances — for example the solution of the one-dimensional Bose gas [2], the six [3] and eight [4] vertex models — brought a totally different approach to the forefront, the Bethe Ansatz. But developments in the last decade have restored the rôle of representation theory. The quantum inverse scattering method [5,6], an algebraic implementation of the Bethe Ansatz and Yang-Baxter equations, gave birth to the theory of quantum groups [7,8]. Representation theory of infinite dimensional Lie algebras became paramount in conformal field theory [9,10].

An independent development was the invention of the corner transfer matrix (CTM) [11,12]. It is the partition function for a whole corner of a regular lattice and might therefore be expected to be even less tractable than the row transfer matrix. Surprisingly, this is not so: CTMs have an extremely simple eigenvalue spectrum (integer powers of a single parameter). CTMs proved to be an effective method for the evaluation of one-point functions [12–14]. In [14] configuration sums appeared which were identified as characters of Virasoro algebras [15,16]. These mysterious connections led to the discovery of hierarchies of solvable lattice models in two dimensions [17] and of many beautiful connections with infinite dimensional Lie algebras [18–20]. Furthermore, the multiplicities of the CTM eigenvalues are equal to the weight-space multiplicities of irreducible highest weight representations of certain affine Lie algebras [20], and the actual configurations used to label the CTM eigenvectors also label the crystal base vectors of the corresponding representations [21–24]. These were vital clues for finding the links with representation theory. But the crucial point was to identify the appropriate algebras and their representations, and the rôle of the row transfer matrices and CTMs in these algebras [25,26].

In one of the original papers on CTMs [11], Baxter observed that there is no Ising-like reduction in the six and eight vertex models. He wrote

A rather ambitious hope is that by examining the CTMs we may stumble on such a group, that the solution of the models may thereby be simplified ...

In [26] was given a new scheme for solving the six-vertex model, in the antiferromagnetic regime, using the newly discovered quantum affine symmetry of the CTM. The approach of that paper has been extended to higher spin chains [27], to the higher rank case [28] and to the ABF models of Andrews, Baxter and Forrester [29]. The scheme has already been used in some of those models to give general expressions for the n -point correlation functions [30]: it also provides a means to derive q -difference equations for them [30,31]. The promise of this new approach is self-evident. Many hard-won results of Bethe Ansatz calculations have been recovered, important new re-

sults have been obtained, and there is much research in progress, since the calculation of correlation functions is a long-standing problem.

The six-vertex model is related to the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ of Drinfeld and Jimbo [8,32], since its Boltzmann weights are the simplest example of an R -matrix. It is a paradigm for the application of the quantum affine symmetry to the representation of CTMs. Briefly, the magic of the six-vertex CTM arises from the property that its eigenvectors provide level-1 highest weight modules of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$, while the CTM acts as a derivation operator in the algebra. Similar statements may be made in the other cases (higher spin, higher rank). Stated more simply, the Chevalley generators of the quantum affine algebra are raising and lowering operators when acting on the eigenvectors of the CTM, and the importance of this cannot be over-stated. It is precisely the kind of property which makes the Ising model so tractable, although the fermion algebra is much simpler than the quantum affine algebra with which we deal here.

The purpose of this article is to give an introductory account which is as elementary as possible. It is based on talks which have been given by the author. For simplicity we consider in detail only the six-vertex model on which the original ideas were developed. In addition, the focus is on the rôle of CTMs; consequently we omit any discussion of the construction and properties of creation/annihilation operators for the XXZ Hamiltonian, although this was a major goal in [26]. There is a growing literature concerned with the extension of the methods to various statistical mechanics systems and with the use of q -difference equations which are an essential ingredient in the theory of q -vertex operators. The six-vertex model uses the level-1 modules of the rank-1 quantum affine algebra $U'_q(\widehat{\mathfrak{sl}}_2)$. Higher-spin and higher-rank cases are considered in [27,28,33]. The Boltzmann weights of all these models form the R -matrices which intertwine tensor products of finite-dimensional representations. The relation is more complicated for the ABF models: the Boltzmann weights are the connection coefficients which intertwine vertex operators, themselves intertwiners [34]. The procedure for the ABF models [29] is somewhat analogous to the Goddard-Kent-Olive coset construction of the discrete minimal series of Virasoro algebras [35]. For the many details of all these developments the reader should consult the cited literature.

2. Transfer matrices

2.1 Definition of the model. The six-vertex model is defined on a regular lattice, each vertex of which is at the intersection of two edges, which carry the physical spin variables $\pm$. The energy E_c of a configuration of the system derives is the sum of the energy E_v assigned to each vertex. From this additivity, the Boltzmann weight $\exp(-E_c/kT)$ of a configuration is the product $\prod_v \exp(-E_v/kT)$ of vertex weights

$$\begin{array}{c} \delta \\ | \\ \alpha \text{ --- } \text{---} \gamma \\ | \\ \beta \end{array} = W(\alpha, \beta, \gamma, \delta; u_{\alpha\gamma} - u_{\beta\delta}).$$

The six-vertex model is “ Z -invariant” in the sense of Baxter [36]. For this reason we attach directions and rapidity variables $u_{\alpha\gamma}$, $u_{\beta\delta}$ to the lines of the lattice, so that $W(\alpha, \beta, \gamma, \delta; u)$ has as its rapidity their difference. The convention is that negating a direction negates the corresponding $u_{\alpha\gamma}$ or $u_{\beta\delta}$.

Only six of the Boltzmann weights are non-zero. In the notation of [37] they are $a = \rho \sinh(\lambda - u)$, $b = \rho \sinh u$, $c = \rho \sinh \lambda$, where ρ is a normalisation factor and $0 \leq u \leq \lambda$ in the physical regime. In anticipation of the connection with the quantum affine algebra, we introduce new variables by

$$q = -\exp(-\lambda), \quad \zeta = \exp(u), \quad \Delta = \frac{q + q^{-1}}{2}.$$

Also, we normalise the weights by the partition function per site, $\kappa(\zeta)$:

$$\begin{aligned} + \begin{array}{c} + \\ | \\ + \end{array} + &= - \begin{array}{c} - \\ | \\ - \end{array} - &= \frac{1}{\kappa(\zeta)}, \\ + \begin{array}{c} - \\ | \\ - \end{array} + &= - \begin{array}{c} + \\ | \\ + \end{array} - &= \frac{1}{\kappa(\zeta)} \frac{1 - \zeta^2}{1 - q^2 \zeta^2} q, \\ + \begin{array}{c} + \\ | \\ - \end{array} - &= - \begin{array}{c} - \\ | \\ + \end{array} + &= \frac{1}{\kappa(\zeta)} \frac{1 - q^2}{1 - q^2 \zeta^2} \zeta. \end{aligned}$$

2.2 Properties of vertex weights. The most important property of the weights is that they satisfy the Yang-Baxter equations, which have the well-known pictorial representation

$$\begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} = \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array}$$

This is the origin of Z -invariance. Summation is implied over all internal lines and this sum is invariant under rearrangement of the lines, provided only that all vertices have the same value of λ and the boundary arrangement is fixed. Because of the locality of the Yang-Baxter equations, this holds for arbitrarily large systems.

Two other important properties are unitarity

$$\begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} = \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array}$$

and crossing symmetry

$$\alpha \begin{array}{c} \delta \\ | \\ \hline | \\ \beta \end{array} \gamma \quad (\zeta) \quad = \quad -\delta \begin{array}{c} \gamma \\ | \\ \hline | \\ \alpha \end{array} -\beta \quad (-q^{-1}\zeta^{-1})$$

The partition function per site may be obtained from these by the standard inversion trick [37]. We sketch the argument. Unitarity and crossing symmetry imply the relations

$$\kappa(\zeta) \kappa(\zeta^{-1}) = 1, \quad \kappa(-q^{-1}\zeta^{-1}) = \frac{1 - q^2 \zeta^2}{q(1 - \zeta^2)} \kappa(\zeta).$$

This gives a q -difference equation and the unique solution for a function analytic in the annulus $q^2 \leq |\zeta| \leq q^{-1}$, and normalised to $\kappa(1) = 1$, is


$$\kappa(\zeta) = \zeta \frac{(\zeta^2 q^4; q^4)_\infty (\zeta^{-2} q^2; q^4)_\infty}{(\zeta^{-2} q^4; q^4)_\infty (\zeta^2 q^2; q^4)_\infty},$$

where

$$(z; p)_\infty = \prod_{n=0}^{\infty} (1 - zp^n).$$

2.3 Corner transfer matrix. CTMs are defined as the partition function of a quadrant of the lattice. The configurations of the edges of the quadrant label the rows and columns of the CTM whilst the outside boundary, which is part of the boundary of the total system, is fixed so as to select the required ground state. There is a preferred diagonal direction (SE – NW), which is along the boundary of $A(u)$, and two ground states labelled $i = 0, 1$, related by a lattice shift of one site.

CTMs are defined in the infinite lattice limit in a normalised form [37]. A small part of the construction of two CTMs, $A(u)$, $B(v)$ looks like:



We consistently use the notation for matrix elements of an operator $L : V \rightarrow W$ that $Lv_\alpha = L_\alpha^\beta w_\beta$: the upper index labels the row.

Recall Baxter's arguments from which the properties of the CTMs stem. He considers the product of two CTMs $A(u)$, $B(v)$. The result is the partition function of an infinite half plane and its value is a matrix element of an arbitrarily high power of the row transfer matrix. This depends only on the maximal eigenvector, which in turn is a function only of the rapidity difference $u - v$ (Yang-Baxter equations). This implies $B(v)A(u) = X(u - v)$. The group property $A(u)A(v) = A(u + v)$ is the most important conclusion to be drawn. It follows from the crossing symmetry of the Boltzmann weights and spin reversal. The result is

$$A(\zeta) = \zeta^{K_{\text{XXZ}}}, \quad B(\zeta) = (-q\zeta)^{-K_{\text{XXZ}}} R,$$

where

$$K_{\text{XXZ}} = \frac{-2}{\sqrt{\Delta^2 - 1}} \sum_{n=1}^{\infty} n(\sigma_{n+1}^x \sigma_n^x + \sigma_{n+1}^y \sigma_n^y + \Delta \sigma_{n+1}^z \sigma_n^z).$$

and R is the operation of spin reversal at every site.

3. The quantum affine algebra

3.1 Notation. In this section we define our notations for $U_q(\widehat{\mathfrak{sl}}_2)$ and review those properties needed for our discussion of the six-vertex model. $U'_q(\widehat{\mathfrak{sl}}_2)$ is generated by e_i , f_i , $t_i = q^{h_i}$, ($i = 0, 1$), which satisfy the defining relations [38]

$$\begin{aligned} t_i e_j &= q^{A_{ij}} e_j t_i, & t_i f_j &= q^{-A_{ij}} f_j t_i, \\ t_i t_j &= t_j t_i, & [e_i, f_j] &= \delta_{ij} \frac{t_i - t_i^{-1}}{q - q^{-1}}, \\ e_i^3 e_j - [3] e_i^2 e_j e_i + [3] e_i e_j e_i^2 - e_j e_i^3 &= 0, \\ f_i^3 f_j - [3] f_i^2 f_j f_i + [3] f_i f_j f_i^2 - f_j f_i^3 &= 0, \end{aligned} \quad (i \neq j).$$

Here $[n] = (q^n - q^{-n})/(q - q^{-1})$, and A_{ij} is the generalised Cartan matrix for the affine Lie algebra $\widehat{\mathfrak{sl}}_2$

$$A_{ij} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}.$$

To obtain the full quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ one must add the generator q^d . d is the derivation, and it satisfies

$$[d, e_i] = \delta_{i,0} e_i, \quad [d, f_i] = -\delta_{i,0} f_i. \quad (1)$$

The co-multiplication for the Hopf algebra structure is defined by

$$\begin{aligned} \Delta(e_i) &= e_i \otimes \text{id} + t_i \otimes e_i, \\ \Delta(f_i) &= f_i \otimes t_i^{-1} + \text{id} \otimes f_i, \\ \Delta(t_i) &= t_i \otimes t_i. \end{aligned}$$

and the formula for the antipode is

$$a(e_i) = -t_i^{-1}e_i, \quad a(f_i) = -f_it_i, \quad a(t_i) = t_i^{-1}.$$

3.2 The R -matrix. The simplest spin half $U_q(\mathfrak{sl}_2)$ module V over the field $\mathbb{Q}(q)$ has basis vectors $v_+ = v_0$, $v_- = v_1$, and is equipped with the actions

$$\begin{aligned} e_1 v_+ &= 0, & f_1 v_+ &= v_-, & t_1 v_+ &= q v_+, \\ e_1 v_- &= v_+, & f_1 v_- &= 0, & t_1 v_- &= q^{-1} v_-. \end{aligned}$$

From this one may construct a $U_q(\widehat{\mathfrak{sl}_2})$ module by introducing the formal variable z . The basis vectors of this infinite-dimensional module are $\{z^n \otimes v_{\pm}\}$ but we simply write $\{z^n v_{\pm}\}$. That is, one equips $V_z = \mathbb{Q}[z, z^{-1}] \otimes V$ with the $U_q(\widehat{\mathfrak{sl}_2})$ module action

$$\begin{aligned} e_0 &\mapsto z f_1, & f_0 &\mapsto z^{-1} e_1, & t_0 &\mapsto t_1^{-1}, & d &\mapsto z \frac{d}{dz}. \\ e_1 &\mapsto e_1, & f_1 &\mapsto f_1, & t_1 &\mapsto t_1, \end{aligned}$$

V_z is also a $U'_q(\widehat{\mathfrak{sl}_2})$ module: for this one simply regards z as a fixed non-zero complex number and drops the grading operator d (evaluation representation).

Define the operator $R(z)$ by

$$\begin{aligned} R(z) v_{\pm} \otimes v_{\pm} &= v_{\pm} \otimes v_{\pm}, \\ R(z) v_+ \otimes v_- &= \frac{1-z}{1-q^2 z} q v_+ \otimes v_- + \frac{1-q^2}{1-q^2 z} z v_- \otimes v_+, \\ R(z) v_- \otimes v_+ &= \frac{1-q^2}{1-q^2 z} v_+ \otimes v_- + \frac{1-z}{1-q^2 z} q v_- \otimes v_+. \end{aligned}$$

Further, let P be the transposition operator: $P \cdot (v_1 \otimes v_2) = (v_2 \otimes v_1)$. Then $\check{R}(z_1/z_2) = PR(z_1/z_2) : V_{z_1} \otimes V_{z_2} \rightarrow V_{z_2} \otimes V_{z_1}$ is an intertwiner of $U'_q(\widehat{\mathfrak{sl}_2})$ modules. Its entries also provide the Boltzmann weights for the six-vertex model with a different normalisation and a change of gauge. Writing $z = \zeta^2$:

$$W(i, j, k, l; \zeta) = \kappa(z)^{-1} g_{ik}^{-1} \check{R}_{ij}^{kl}(z),$$

with $g_{ik} = 1$, $i = k$, and $g_{\pm\mp} = \zeta^{\pm 1}$. The gauge factor is the well-known term which makes finite spin chains $U_q(\mathfrak{sl}_2)$ -invariant. It simply adds the total spin to K_{XZX} and changes its rôle as a grading operator. Since it does not affect the physics, we henceforth use the Boltzmann weights

$$\overline{W}(i, j, k, l; z) = g_{ik} W(i, j, k, l; \zeta).$$

This is consistent with the cited works [26–29, 33].

3.3 Dual modules. The dual module construction is intimately connected with the crossing symmetry and is required for representing local operators. It goes as follows.

Given a linear space V its dual V^* is the set of linear maps $V^* : V \rightarrow \mathbb{C}$ by the action $v \xrightarrow{u^*} \langle u^*, v \rangle$. Choose the dual basis by the canonical pairing $\langle v_j^*, v_k \rangle = \delta_{jk}$. V_z^* becomes a dual module under the action

$$\langle xu^*, v \rangle = \langle u^*, \phi(x)v \rangle, \quad \forall x \in U_q(\widehat{\mathfrak{sl}_2}),$$

where ϕ is any anti-homomorphism of the algebra ($\phi(xy) = \phi(y)\phi(x)$). Using the antipode for ϕ , the action of $U_q(\widehat{\mathfrak{sl}_2})$ on V_z^* is

$$\begin{aligned} e_1 v_+^* &= -q^{-1} v_-^*, & f_1 v_+^* &= 0, & t_1 v_+^* &= q^{-1} v_+^*, \\ e_1 v_-^* &= 0, & f_1 v_-^* &= -q v_-^*, & t_1 v_-^* &= q v_-^*, \\ e_0 &\mapsto (zq^{-2}) f_1, & f_0 &\mapsto (zq^{-2})^{-1} e_1, & t_0 &\mapsto t_1^{-1}. \end{aligned}$$

This gives the isomorphism

$$C : V_z^* \xrightarrow{\sim} V_{zq^{-2}}, \quad v_+^* \mapsto v_-, \quad v_-^* \mapsto -q v_+.$$

The operator $R(z)$ was defined to act on $V_{z_1} \otimes V_{z_2}$ so that $PR(z)$ is an intertwiner. We also require the equivalent operator $R^*(z)$ acting on $V_{z_1}^* \otimes V_{z_2}$. It is given as $R^*(z) = (C \otimes \text{id})^{-1} R(zq^{-2})(C \otimes \text{id})$. Explicitly,

$$\begin{aligned} R^*(z) v_\pm^* \otimes v_\mp &= v_\pm^* \otimes v_\mp, \\ R^*(z) v_+^* \otimes v_+ &= \frac{q - q^{-1}z}{1 - z} v_+^* \otimes v_+ + \frac{q - q^{-1}}{1 - z} v_-^* \otimes v_-, \\ R^*(z) v_-^* \otimes v_- &= \frac{q - q^{-1}}{1 - z} z v_+^* \otimes v_+ + \frac{q - q^{-1}z}{1 - z} v_-^* \otimes v_-. \end{aligned}$$

3.4 Notation. Our notations for $\widehat{\mathfrak{sl}_2}$ are as follows. The Cartan subalgebra $\mathfrak{H}$ is spanned by $\{h_0, h_1, d\}$ and α_0, α_1 are the roots. They are related to the fundamental weights by $\alpha_0 = 2\Lambda_0 - 2\Lambda_1 + \delta$, $\alpha_1 = 2\Lambda_1 - 2\Lambda_0$; we also write $\rho = \Lambda_0 + \Lambda_1$. The invariant form on $\mathfrak{H}^*$ is given by $2(\Lambda_i, \Lambda_j) = \delta_{1i}\delta_{1j}$, $(\Lambda_i, \delta) = 1$, $(\delta, \delta) = 0$. The weight lattice and its dual are $P = \mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\Lambda_1 \oplus \mathbb{Z}\delta$ and $P^* = \mathbb{Z}h_0 \oplus \mathbb{Z}h_1 \oplus \mathbb{Z}d$, with $\langle \Lambda_i, h_j \rangle = \delta_{ij}$, $\langle \Lambda_i, d \rangle = 0$, $\langle \delta, h_i \rangle = 0$, $\langle \delta, d \rangle = 1$. We identify P^* with a subset of P via $(\ , \)$, so that $\alpha_i = h_i$ and

$$2\rho = 4d + h_1. \tag{2}$$

This plays an important rôle in relating K_{XXZ} and K_{CTM} with ρ and d .

There are two irreducible infinite dimensional level 1 highest weight $U_q(\widehat{\mathfrak{sl}_2})$ modules, $V(\Lambda_0)$, $V(\Lambda_1)$, corresponding to the fundamental weights Λ_0, Λ_1 . Their relation to the eigenvectors of the CTM will be discussed in the next section.

4. The CTM and its eigenvectors

In [25] Foda and Miwa identified a six-vertex CTM generator with the derivation d of $U_q(\widehat{\mathfrak{sl}_2})$, and its eigenvectors with the weight vectors of the level-1 modules. There are two bare ground states, $(\cdots, -, +, -, +)$ and $(\cdots, +, -, +, -)$. They defined the action of the algebra on the sets of paths $\mathcal{P}_i$, $(i = 0, 1)$, which differ from these states at only a finite number of places. The problem is to show that everything works consistently in each order of perturbation. Convincing evidence was given for this, through extensive analytic computations to quite high powers of q , but the main result remained a conjecture. Moreover the q -expansions about these bare states are divergent, and an infinite renormalisation is involved.

In the next two subsections we explain briefly the approach of Davies [39], which uses finite size truncation together with the Kashiwara's crystal base theory [21,22] to prove the necessary identifications. This method generalises quite easily to the higher level and higher rank cases.

4.1 Identification of derivation. K_{XXZ} is defined using the Boltzmann weights W , but it is convenient to introduce another generator K_{CTM} defined from $\overline{W}$. It is built from two-site operators H_j :

$$H_j = \frac{1}{2(q - q^{-1})} \left(\sigma_{j+1}^x \sigma_j^x + \sigma_{j+1}^y \sigma_j^y + \Delta \sigma_{j+1}^z \sigma_j^z + \frac{(q - q^{-1})}{2} (\sigma_{j+1}^z - \sigma_j^z) \right).$$

obtained by expansion of $\check{R}$ about $z = 1$. The subscript j indicates that H_j operates at the positions $j + 1, j$ of the chain. For this reason, we attach a second subscript to the generators e_i, f_i, t_i , writing $e_{i,j}, f_{i,j}, t_{i,j}$. The truncated CTM generator is

$$K_{\text{CTM}}^{(N)} = \sum_{j=1}^{N-1} j H_j.$$

It acts on the N -fold tensor product $V^{\otimes N}$. Write $\hat{e}_i, \hat{f}_i, \hat{t}_i$ for the N -fold iterated co-product of the generators,

$$\begin{aligned} \hat{e}_i &= \sum_{j=1}^N t_{i,N} \otimes \cdots \otimes t_{i,j+1} \otimes e_{i,j} \otimes \text{id}_{j-1} \otimes \cdots \otimes \text{id}_1, \\ \hat{f}_i &= \sum_{j=1}^N \text{id}_N \otimes \cdots \otimes \text{id}_{j+1} \otimes f_{i,j} \otimes t_{i,j-1}^{-1} \otimes \cdots \otimes t_{i,1}^{-1}, \\ \hat{t}_i &= t_{i,N} \otimes \cdots \otimes t_{i,1}. \end{aligned}$$

We need their commutation relations with $K_{\text{CTM}}^{(N)}$. For $\hat{e}_1, \hat{f}_1$ this is trivial, $\check{R}(x, y)$ intertwines the action of the subalgebra $U_q(\mathfrak{sl}_2)$ which does not depend on the spectral parameter. It follows immediately that $K_{\text{CTM}}^{(N)}$ is $U_q(\mathfrak{sl}_2)$ invariant:

$$[K_{\text{CTM}}^{(N)}, \hat{e}_1] = 0, \quad [K_{\text{CTM}}^{(N)}, \hat{f}_1] = 0.$$

The intertwining condition for e_0, f_0 gives the commutation relations [39]

$$\begin{aligned} [K_{\text{CTM}}^{(N)}, \hat{e}_0] &= \hat{e}_0 - N e_{0,N} \otimes \text{id}_{N-1} \cdots \otimes \text{id}_1, \\ [K_{\text{CTM}}^{(N)}, \hat{f}_0] &= -\hat{f}_0 + N f_{0,N} \otimes t_{0,N-1}^{-1} \otimes \cdots \otimes t_{0,1}^{-1}. \end{aligned}$$

This is the vital result. In the infinite N limit the boundary terms may be neglected, so that $K_{\text{CTM}} = \lim_{N \rightarrow \infty} (K_{\text{CTM}}^{(N)} - R_N)$ acts as the derivation in the algebra $U_q(\widehat{\mathfrak{sl}}_2)$, verifying the relations (1). R_N is a renormalisation constant and plays no further rôle.

4.2 Identification of modules. K_{XXZ} and K_{CTM} act formally on a semi-infinite tensor product of spin-half modules, as does the quantum affine algebra. It must be emphasised that this limit is only formally defined. When the eigenstates are developed either as ℓ_2 -sequences or as q -expansions, one obtains divergent expressions. The situation is similar to the Ising model CTMs [40,41]. Having truncated the chain at a finite length, the action of the algebra is well-defined. $K_{\text{CTM}}^{(N)}$ is renormalised by its ground state eigenvalue. It is shown in [39] that the eigenvectors with eigenvalues up to M carry the $U_q(\widehat{\mathfrak{sl}}_2)$ action to within error terms of order $q^{2N-2M-1}$. So the boundary terms carry factors of the order $Nq^{\approx 2N}$, which is why they may be neglected in the infinite limit. It is also shown there, using the theory of the crystal base, that the eigenvectors are in one-to-one correspondence with the weight vectors of the modules $V(\Lambda_i)$.

This method of finite size approximation had already been noted for the crystal base [24], so it is natural to extend it to $q \neq 0$. The meaning of the approximation is quite precise, and illustrates the non-uniform nature of the convergence. For any fixed integer M , the eigenstates corresponding to the grading levels $\leq M$ approximate weight vectors up to that grading level to within errors of order $Nq^{2N-4M-1}$ as $N \rightarrow \infty$. But if one selects some order of approximation, and then increases M and N together, the approximated eigenvectors become an increasingly small fraction of the totality of eigenvectors of the truncated CTM, since $2^N/2^M$ increases exponentially fast. What matters is that q -expansions of physical quantities obtained from the CTM and from representation theory will agree to every order in q .

Without further ado, we shall identify the eigenvectors of the CTM with the weight vectors of the highest weight modules, and the generator K_{CTM} with the derivation operator d . Examination of the gauge term in $\overline{W}$ shows that $K_{\text{XXZ}} = 2K_{\text{CTM}} + S$, $2S = \sum_{k=1}^{\infty} \sigma_j^z$. Using the connections (2) leads to the identification

$$K_{\text{XXZ}} = 2d + h_1/2 = \rho.$$

4.3 Local properties. Let L be a local operator, acting at the sites $1, \dots, n$. The expectation value $\langle L \rangle$ is calculated by the standard CTM method [37] as a quotient of traces:

$$\langle L \rangle = \frac{\text{tr}(LABCD)}{\text{tr}(ABCD)} = \frac{\text{tr}_{V(\Lambda_i)}(Lq^{2\rho})}{\text{tr}_{V(\Lambda_i)}(q^{2\rho})}. \quad (3)$$

Here A , B , C and D are CTMs for the four quadrants of the system. The difficulty is that L must be expressed in a basis of eigenvectors of K_{XXZ} . Without an explicit construction for these, the calculation is restricted to an operator which commutes with K_{XXZ} , i.e., a one-point function. For the six-vertex model this $L = (-1)^S$ and the one-point function is the magnetisation in the “interaction round a face” representation [37]. The traces are specialisations of the character formulae:

$$\chi_i(x, y) = \text{tr}_{V(\Lambda_i)}(x^d y^{h_1}) = y^{-i} \frac{(-x^{1-i} y^2; x^2)_\infty (-x^{1+i} y^{-2}; x^2)_\infty}{(x; x^2)_\infty}.$$

Substituting $x = q^4$, $y = q$ into these formulae gives the magnetisation as

$$m = \frac{(q^2; q^4)_\infty}{(-q^2; q^4)_\infty} = \prod_{n=0}^{\infty} \frac{(1 - q^{4n+2})}{(1 + q^{4n+2})}.$$

5. Local description of CTM eigenvectors

5.1 Change of basis. The difficulty of defining the CTM eigenvectors as normalisable ℓ_2 sequences has already been noted. Representation theory provides a way around the problem. Consider the tensor product $V(\Lambda_{1-i}) \otimes V$. Formally, we identify this with $(\otimes_{n=2}^{\infty} V_n) \otimes V_1$, with boundary condition for sector i , but counted from the first site. (There is some abuse of notation: V_n here denotes the spin-half module for site n .) The product may also be written $(\otimes_{n=1}^{\infty} V_n)$ and then equated to weight vectors of $V(\Lambda_i)$. The map between the two pictures is a q -vertex operator (type I Vertex Operators or VO [34]) of Frenkel and Reshetikhin [34].

$$\dots \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \xrightarrow{\sim} \quad \dots \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array} \quad \otimes \quad \begin{array}{|c|} \hline \text{---} \\ \hline \end{array}$$

In this diagram crosses represents sites in the chain (copies of V), not vertices in transfer matrices.

VOs are simply homomorphisms — they preserve the algebra structure. They may also be interpreted as expansions of the CTM eigenvectors for boundary sector i in terms of the state variable at site 1 and the eigenvectors for the boundary sector $1 - i$. This process may be iterated to obtain information about any finite number of state variables at sites $1, \dots, n$.

VOs give precisely the information required for the calculation of local properties in (3). Moreover, the expansions have convergent domains in q and z . But the most crucial property is that the intertwining property of the VOs uniquely determines them to within normalisation. Once one has made the identifications of section 4, no further physical arguments are required. The introductory comments above, and those of section 5.3 below, are quite superfluous in this sense. They are included to provide some intuitive understanding of the physical meaning of the theory.

5.2 Vertex operators. The basic definition of VOs is that they are $U'_q(\widehat{\mathfrak{sl}}_2)$ homomorphisms

$$\begin{aligned} \Phi_{\Lambda_i}^{\Lambda_{1-i}V} &: V(\Lambda_i) \rightarrow V(\Lambda_{1-i}) \otimes V, \\ \Phi_{\Lambda_{1-i}V}^{\Lambda_i} &: V(\Lambda_{1-i}) \otimes V \rightarrow V(\Lambda_i). \end{aligned}$$

We discuss $\Phi_{\Lambda_i}^{\Lambda_{1-i}V}$. It gives an expansion of weight vectors $u \in V(\Lambda_i)$ as tensor products:

$$u = \sum_{\substack{n \in \mathbb{Z} \\ j = \pm}} w_{n,j} \otimes v_j, \quad w_{n,j} \in V(\Lambda_{1-i}).$$

The intertwining property

$$\Phi_{\Lambda_i}^{\Lambda_{1-i}V}(xu) = \Delta(x) \Phi_{\Lambda_i}^{\Lambda_{1-i}V}u, \quad \forall x \in U'_q(\widehat{\mathfrak{sl}_2}),$$

implies that $\text{wt}(u) = \text{wt}(w_{n,j}) + \text{wt}(v_j) + n\delta$. Plainly speaking, the expansion is over states of all energy whose spins match. The normalisation is fixed by setting the coefficient of the term in the highest weight vectors u_{Λ_i} to unity:

$$\begin{aligned} \Phi_{\Lambda_i}^{\Lambda_{1-i}V}u_{\Lambda_i} &= (u_{\Lambda_{1-i}} \otimes v_{1-i}) + \cdots, \\ \Phi_{\Lambda_{1-i}V}^{\Lambda_i}(u_{\Lambda_{1-i}} \otimes v_{1-i}) &= u_{\Lambda_i} + \cdots. \end{aligned}$$

VOs may also be defined to preserve the grading operator d : they are the $U_q(\widehat{\mathfrak{sl}_2})$ homomorphisms

$$\begin{aligned} \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z) : V(\Lambda_i) &\rightarrow V(\Lambda_{1-i}) \otimes V_z, \\ \tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z) : V(\Lambda_{1-i}) \otimes V_z &\rightarrow V(\Lambda_i). \end{aligned}$$

The two forms (Φ and $\tilde{\Phi}(z)$) are trivially related by inserting or deleting the appropriate powers of z ,

$$u = \sum_{\substack{n \in \mathbb{Z} \\ j = \pm}} w_{n,j} \otimes z^n v_j.$$

Since the derivation detects the grading before and after the action of the VO, the powers of z may be inserted by conjugation with the operator z^d :

$$\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z) = z^{d \otimes \text{id}} \Phi_{\Lambda_i}^{\Lambda_{1-i}V} z^{-d}.$$

From this follows the “homogeneity property”

$$\begin{aligned} \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z_1 z_2) &= z_1^{d \otimes \text{id}} \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z_2) z_1^{-d}, \\ \tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z_1 z_2) &= z_1^d \tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z_2) z_1^{-d \otimes \text{id}}. \end{aligned}$$

Some elementary calculations give the following results:

$$\begin{aligned} \tilde{\Phi}_{\Lambda_0}^{\Lambda_1 V}(z) u_{\Lambda_0} &= u_{\Lambda_1} \otimes v_- - q(f_1 u_{\Lambda_1}) \otimes v_+ + \mathcal{O}(z), \\ \tilde{\Phi}_{\Lambda_1}^{\Lambda_0 V}(z) u_{\Lambda_1} &= u_{\Lambda_0} \otimes v_+ + \mathcal{O}(z), \\ \tilde{\Phi}_{\Lambda_1}^{\Lambda_0 V}(z)(f_1 u_{\Lambda_1}) &= u_{\Lambda_0} \otimes v_- + \mathcal{O}(z). \end{aligned} \tag{4}$$

They will be used below to fix normalisations.

5.3 *Physical description.* Let $P_{\beta}^{\gamma,j}$ be a non normalised semi-infinite shift operator with entries

$$P_{\dots,\beta_3,\beta_2,\beta_1}^{\dots,\gamma_2,\gamma_1,j} \simeq \dots \delta_{\beta_3,\gamma_2} \delta_{\beta_2,\gamma_1} \delta_{\beta_1,j}.$$

Since P preserves the $U'_q(\widehat{\mathfrak{sl}}_2)$ action one may identify an appropriate constant multiple with the VO $\Phi_{\Lambda_i}^{\Lambda_{1-i}V}$. Further, we use the relation of d with the CTM to write $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z) = A_{1-i}(z)PA_i(z^{-1})$. We may even identify the VO as a normalised row transfer matrix. This is a generalisation of the boost property for the full transfer matrix [42,43]. The product PA_i is the partition function of a quadrant with the boundary type i along its horizontal edge and $1-i$ along its vertical one. It is not a CTM, but may be re-written as a different product:

The semi-infinite row of vertices corresponds to the vertex operator $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z)$, with a normalisation factor $\phi_i(z)$.

This physical interpretation of VOs follows [31].

The product of row transfer matrices is commuted using the Yang Baxter equations together with the assumption that boundary effects disappear for a system with fixed boundary conditions:

We use this to fix normalisation factors. Write

$$\tilde{R}(z_1/z_2) \tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_1) \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2) = \kappa(z_1/z_2) \frac{\phi_{1-i}(z_2)\phi_i(z_1)}{\phi_{1-i}(z_1)\phi_i(z_2)} \tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_2) \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_1),$$

and take the vacuum expectation. From the definitions, the terms in $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z_1)u_{\Lambda_i}$ involve only non-negative powers of z_1 . In order to arrive once more at the highest weight vector after the action of $\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_2)$, these terms must pair with a matching negative power of z_2 . Denote the vacuum expectation as

$$\Psi_i(z_1/z_2) = \langle u_{\Lambda_i} | \tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_1) \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2) | u_{\Lambda_i} \rangle.$$

It involves only non-negative powers of z_1/z_2 , and will be analytic in z_1/z_2 in some domain containing the origin. As an element of $V \otimes V$ it must be an eigenvector of $\check{R}(z_1/z_2)$ because of the commutation relation. A simple calculation gives:

$$\begin{aligned}\check{R}(z)(v_+ \otimes v_- - qv_- \otimes v_+) &= \frac{z - q^2}{1 - q^2 z} (v_+ \otimes v_- - qv_- \otimes v_+), \\ \check{R}(z)(v_- \otimes v_+ - qz^{-1}v_+ \otimes v_-) &= \frac{1 - q^2 z^{-1}}{1 - q^2 z} (v_- \otimes v_+ - qz^{-1}v_+ \otimes v_-).\end{aligned}$$

To leading order this corresponds exactly with (4). Inserting the formula for $\kappa(z)$ and using analyticity,

$$\begin{aligned}\Psi_0(z) &= \frac{(q^6 z; q^4)_\infty}{(q^4 z; q^4)_\infty} (v_+ \otimes v_- - qv_- \otimes v_+), \\ \Psi_1(z) &= \frac{(q^6 z; q^4)_\infty}{(q^4 z; q^4)_\infty} (v_- \otimes v_+ - qz^{-1}v_+ \otimes v_-).\end{aligned}\tag{5}$$

and

$$\phi_i(z) = \gamma_i z^{\Delta_{1-i} - \Delta_i}, \quad \Delta_0 = 0, \quad \Delta_1 = 1/4,$$

for some constants γ_i . The “conformal factors” Δ_i , which were included in the definition of the VOs in [26], are a shift in the zero point; $A_i(z) \simeq z^{d+\Delta_i}$. Such shifts are calculated for the Ising model in [41].

5.4 Mathematical construction. We have discussed the physical construction of the VOs in some detail. However, for the purpose of calculation, the mathematical construction is preferable. The scale factors (5) are determined unambiguously in [26,28] by solving the q -KZ (Knihnik-Zamolodchikov) equation [34]. From this, the commutation relation for the VOs follows immediately, viz:

$$\begin{aligned}PR(z_1/z_2)\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_1)\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2) &= z^{\delta_{i0}} r(z_1/z_2)\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_2)\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_1), \\ r(z) &= \frac{(zq^4; q^4)_\infty (z^{-1}q^6; q^4)_\infty}{(z^{-1}q^4; q^4)_\infty (zq^6; q^4)_\infty}.\end{aligned}$$

This confirms our identification of the conformal factors, and the rôle of the partition function, in interpreting the VOs as transfer matrices.

The charge conjugation isomorphism C relates $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V^*}(z)$ to $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z)$ by

$$\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V^*}(z) = (-q)^{\delta_{i1}} (\text{id} \otimes C)^{-1} \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(zq^{-2}),$$

and from this the commutation relation emerges in a form needed for the calculation of the spontaneous polarisation. It is

$$PR^*(z)\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V^*}(z_1)\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2) = -q^{-1} z^{\delta_{i0}} r(zq^{-2})\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_2)\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V^*}(z_1).$$

To complete the determination of local operators, we must construct the inverse of $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z)$. By definition, it is $\tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z)$ up to normalisation. The construction goes as follows. The VOs are algebra homomorphisms, so one may use the canonical identification that $\text{Hom}(L \otimes N, M)$ is isomorphic to $\text{Hom}(L, M \otimes N^*)$, to equate $\tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z)$ with $\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V^*}(z)$. Let us expand on this. It means that

$$\tilde{\Phi}_{\Lambda_{1-i}V}^{\Lambda_i}(z)(u \otimes v_j) = \langle \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V^*}(z)u, v_j \rangle,$$

where the pairing is in $V^* \otimes V$. One may check that this is correct on the Chevalley generators. For example,

$$\begin{aligned} \tilde{\Phi}_{\Lambda_i V}^{\Lambda_{1-i}}(z)(\Delta(e_i)(u \otimes v_j)) &= \langle e_i \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V^*}(z)u, v_j \rangle \\ &+ \sum_k \left(t_i \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V^*}(z)u \right)_k \left(\langle e_i v_k^*, v_j \rangle + \langle t_i v_k^*, e_i v_j \rangle \right). \end{aligned}$$

The second line vanishes because the antipode is used to define the dual. Finally, (5) tells us the scalar factor for the inverse. Setting $z = q^{-2}$ gives

$$\Phi_{\Lambda_{1-i}V}^{\Lambda_i}(z)\Phi_{\Lambda_i}^{\Lambda_{1-i}V}(z) = \frac{(q^4; q^4)_\infty}{(q^2; q^4)_\infty} \text{id}_{V(\Lambda_i)}.$$

This completes the construction of change of basis matrices.

5.5 Spontaneous polarisation. Two decades ago, Baxter calculated the spontaneous polarisation of the six-vertex model using the Bethe Ansatz [44]. Unlike many other important results, no alternative derivation was subsequently forthcoming, although the method has recently been extended to the calculation of the mean polarisation in the higher spin case [45]. The properties of VOs gives the first new derivation: moreover VOs give a general method to obtain correlation functions. The ideas have even been used for the eight-vertex model [46], validating a long-standing conjecture [47], although the underlying algebraic structure is not presently understood in that case.

Here we sketch the derivation for the six-vertex model, which is rank-1. The solution for arbitrary rank has recently been given by Koyama [33]. The expectation value that a single edge variable has the value m ($= \pm$) is simply the trace of an operator with matrix elements $E_{jk} = \delta_{jm}\delta_{km}$. Recalling our identification of the CTM generator K_{CTM} with ρ , consider the more general unnormalised expectation value

$$F^{(i)}(z_1/z_2) = \text{tr}_{V(\Lambda_i)}(q^{-2\rho} \tilde{\Phi}_{\Lambda_{i-1}}^{\Lambda_i V^*}(z_1) \tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i}V}(z_2)).$$

It takes its values in $V^* \otimes V$ and the spontaneous polarisation is the ratio $(F_{++}^{(i)}(1) - F_{--}^{(i)}(1))/(F_{++}^{(i)}(1) + F_{--}^{(i)}(1))$.

We may derive a q -difference equation for $F^{(i)}$:

$$\begin{aligned}
F^{(i)}(z_1/z_2q^4) &= \text{tr}_{V(\Lambda_i)}(q^{-2\rho}\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V^*}(z_1)\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2q^4)) \\
&= P \text{tr}_{V(\Lambda_{1-i})}(\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2q^4)q^{-2\rho}\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V^*}(z_1)) \\
&= P(q^{-h_1} \otimes \text{id}) \times \text{tr}_{V(\Lambda_{1-i})}(q^{-2\rho}\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V}(z_2)\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V^*}(z_1)) \\
&= -qz^{-\delta_{i1}}r(zq^{-2})(\text{id} \otimes q^{-h_1})R^*(z) \times \text{tr}_{V(\Lambda_{1-i})}(q^{-2\rho}\tilde{\Phi}_{\Lambda_i}^{\Lambda_{1-i} V^*}(z_1)\tilde{\Phi}_{\Lambda_{1-i}}^{\Lambda_i V}(z_2)).
\end{aligned}$$

The three steps use, respectively, the cyclic property of the trace, the homogeneity and intertwining property of the VOs, and the commutation relations. The equation thus obtained reads

$$F^{(i)}(zq^{-4}) = -qz^{-\delta_{i1}}(\text{id} \otimes q^{-h_1})r(zq^{-2})R^*(z)F^{(1-i)}(z).$$

These equations may be solved quite simply. Set $G^{(\pm)}(\zeta) = F^{(1)}(\zeta^2) \pm q\zeta F^{(0)}(\zeta^2)$, which decouples them. Note also that only the 2 by 2 block spanned by $v_{\pm}^* \otimes v_{\pm}$ is non zero. For it we find

$$\begin{aligned}
G^{(\pm)}(\zeta q^{-2}) &= \mp r(\zeta^2 q^{-2})M(\zeta)G^{(\pm)}(\zeta), \\
M(\zeta) &= \frac{\zeta}{1-\zeta^2} \begin{pmatrix} 1-q^{-2}\zeta^2 & \zeta^2(1-q^{-2}) \\ q^2-1 & q^2-\zeta^2 \end{pmatrix}.
\end{aligned}$$

This may be reduced to scalar equations by considering a generalised eigenvalue problem: $M(\zeta)w(\zeta) = \lambda(\zeta)w(\zeta q^{-2})$. The eigenvectors are

$$\frac{w_{\pm}(\zeta)}{(1 \mp \zeta)} = (1, \pm \zeta^{-1}), \quad \lambda(\zeta) = \zeta.$$

The solution for $G^{(\pm)}(\zeta)$ involves $\Theta_{q^2}(\pm\zeta)$, where

$$\Theta_p(\zeta) = (p; p)_{\infty}(\zeta; p)_{\infty}(\zeta^{-1}p; p)_{\infty}.$$

It satisfies $\Theta_p(\zeta p) = -\zeta^{-1}\Theta_p(\zeta)$. In order to have the required analyticity a zero of this function must cancel the factor $(1 \mp \zeta)$ in $w_{\pm}(\zeta)$ and this implies that there is only one non-zero component for each of $G^{(\pm)}(\zeta)$. The argument whereby one shows that $F^{(i)}(z)$ has the assumed analyticity is based on finding an integral representation, and is given in [33,46]. The solutions may be normalised by the fact that $F^{(i)}(q^{-2})$ is the trace of the identity operator. However, this is not necessary for finding the staggered polarisation. The $G^{(\pm)}(\zeta)$ differ only in the factors $\Theta_{q^2}(\pm\zeta)$, and we find, independent of i ,

$$P = \left[\frac{(1+\zeta)\Theta_{q^2}(\zeta)}{(1-\zeta)\Theta_{q^2}(-\zeta)} \right]_{\zeta=1} = \prod_{n=0}^{\infty} \frac{(1-q^{2n+2})^2}{(1+q^{2n+2})^2}.$$

The beautiful interplay which has emerged between physics and mathematics brings to mind a quote from the writings of Ludwig Boltzmann:


It is unbelievable how simple and straight forward each result appears once it has been found, and how difficult it seems so long as the way which leads to it is unknown.

Acknowledgements

I wish to acknowledge many helpful discussions with D. Altschuler, M. T. Batchelor, R. J. Baxter, A. Carey, O. Foda, M. Jimbo, T. Miwa, M. Okado, P. A. Pearce, I. Peschel, N. Yu. Reshetikhin, R. W. Richardson and R. Zhang.


Appendix: Some numerical results

For making finite size numerical calculations it is convenient to use $K_{\text{CTM}}^{(N)}$ rather than $K_{\text{XXZ}}^{(N)}$ because the gauge term forces the boundary conditions to those appropriate for $V(\Lambda_0)$ when N is even and $V(\Lambda_1)$ when N is odd [39]. Here we show some results for chains of length up to 11. Energies are relative to the antiferromagnetic ground state energy. They should be compared with the figures for the weight space structure of the relevant modules. $U_q(\mathfrak{sl}_2)$ -invariance means that the $\hat{f}_1$ multiplets are exact. The $\hat{f}_0$ multiplets tend to the proper degeneracy as N increases. Numerical checks have also been made on the actions of $\hat{e}_0$ and $\hat{f}_0$ for these values of N . In the weight space diagrams for the level-1 modules, spin is in the horizontal direction and grading is downwards. The action of f_1 is shown as a left pointing arrow, since f_1 decreases the weight by α_1 . f_0 is a diagonal arrow down and to the right, since f_0 decreases the weight by α_0 , and $\alpha_0 + \alpha_1 = \delta$.



Spin	$N = 4$	$N = 6$	$N = 8$	$N = 10$
0	0	0	0	0
1	1.0021201	1.0002473	1.0000301	1.0000038
0	2.1189021	2.0203449	2.0033092	2.0005036
1	2.1167936	2.0200996	2.0032794	2.0004998
0	—	3.2770532	3.0678881	3.0151265
1	3.1200455	3.0205449	3.0033418	3.0005085
1	—	3.2768064	3.0678580	3.0151227
0	—	4.2968794	4.0711308	4.0156211
0	—	—	4.4835212	4.1520062
1	—	4.2771017	4.2771017	4.0151267
1	—	—	4.4839116	4.1520024
2	—	4.0205469	4.0033421	4.0005086

Weight spaces of $V(\Lambda_0)$ and negative eigenvalues of truncated CTM



Spin	$N = 5$	$N = 7$	$N = 9$	$N = 11$
1/2	0	0	0	0
1/2	1.0081308	1.0011185	1.0001513	1.0000204
1/2	2.1899754	2.0396425	2.0076092	2.0013141
3/2	2.0082121	2.0011297	2.0001528	2.0000206
1/2	3.1980888	3.0407572	3.0077598	3.0013343
1/2	—	3.3752382	3.1052883	3.0268467
3/2	3.1899870	3.0396431	3.0076093	3.0013141
1/2	4.2000343	4.0411612	4.0078363	4.0013475
1/2	—	4.3763556	4.1054395	4.0268671
1/2	—	—	4.6000763	4.2077221
3/2	4.1999428	4.0411497	4.0078347	4.0013473
3/2	—	4.3752396	4.1052884	4.0268467

Weight spaces of $V(\Lambda_1)$ and negative eigenvalues of truncated CTM

References

- [1] L. Onsager: Crystal statistics. I. A two-dimensional model with an order-disorder transition, *Phys. Rev.*, **65**, (1944), 117-149.
- [2] E. H. Lieb and W. Liniger: Exact analysis of an interacting Bose gas: I. The general solution and the ground state, *Phys. Rev.*, **130**, (1963), 1605-1616.
— : Exact analysis of an interacting Bose gas: II. The excitation spectrum, *Phys. Rev.*, **130**, (1963), 1616-1624.
- [3] E. H. Lieb: Residual entropy of square ice, *Phys. Rev.*, **162**, (1967), 162-172.
— : Exact solution of the F model of an antiferroelectric, *Phys. Rev. Lett.*, **18**, (1967), 1046-1048.
— : Exact solution of the two dimensional Slater KDP model of a ferroelectric, *Phys. Rev. Lett.*, **19**, (1967), 108-110.
B. Sutherland: Exact solution of a two dimensional model for Hydrogen bonded crystals, *Phys. Rev. Lett.*, **19**, (1967), 103-104.
- [4] R. J. Baxter: Partition function of the eight-vertex lattice model, *Ann. Phys.*, NY, **70**, (1972), 193-228.
- [5] L. D. Faddeev: Quantum completely integrable models in field theory, *Sov. Sci. Rev. Math. Phys.*, **C1**, (1980), 107-155.
- [6] H. B. Thacker: Exact integrability in quantum field theory and statistical systems, *Rev. Mod. Phys.*, **53**, (1981), 253-285.
- [7] M. Jimbo: A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation, *Lett. Math. Phys.*, **10**, (1985), 63-69.
- [8] V.G. Drinfeld, *Quantum groups*, *Proc. ICM Berkeley*, (1986), 798-820.
- [9] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov: Infinite conformal symmetry in two dimensional quantum field theory, *Nucl. Phys. B*, **241**, (1984), 333-380.

- [10] G. Felder and A. LeClair: Restricted quantum affine symmetry of restricted minimal conformal models, *Int. J. Mod. Phys. A*, **7**, Suppl. 1A, (1992), 239-278.
- [11] R. J. Baxter: Corner transfer matrices of the eight vertex model. Low temperature expansions and conjectured properties, *J. Stat. Phys.*, **15**, (1976), 485-503.
- [12] R. J. Baxter : Corner transfer matrices, *Physica A*, **106**, (1981), 18-27.
- [13] R. J. Baxter: Hard Hexagons: exact solution, *J. Phys. A: Math. Gen.*, **13**, (1980), L61-L70.
— : Rogers-Ramanujan identities in the hard hexagon model, *J. Stat. Phys.*, **26**, (1981), 427-452.
- [14] G. E. Andrews, R. J. Baxter and P. J. Forrester: Eight-vertex SOS model and generalized Rogers-Ramanujan type identities, *J. Stat. Phys.*, **35**, (1984), 193-266.
- [15] J. L. Cardy: Conformal invariance and surface critical behaviour, *Nucl. Phys. B*, **240**, (1984), 514-532.
- [16] D. Friedan, Z. Qiu and S. Shenker, *Vertex operators in Mathematical Physics*, eds. J. Lepowski, S. Mandelstam and I. M. Singer, Springer (1984).
- [17] M. Jimbo, T. Miwa and M. Okado: Solvable lattice models related to the vector representation of classical simple Lie algebras, *Comm. Math. Phys.*, **116**, (1988), 507-525.
- [18] E. Date, M. Jimbo, A. Kuniba, T. Miwa and M. Okado: Exactly solvable SOS models: Local height probabilities and theta function identities, *Nucl. Phys. B*, **290**, (1987), 231-273.
- [19] M. Jimbo, T. Miwa and M. Okado: Local state probabilities of solvable lattice models: An $A_{n-1}^{(1)}$ family, *Nucl. Phys. B*, **300**, (1988), 74-108.
- [20] E. Date, M. Jimbo, A. Kuniba, T. Miwa and M. Okado: One dimensional configuration sums in vertex models and affine Lie algebra characters, *Lett. Math. Phys.*, **17**, (1987), 69-77.
- [21] M. Kashiwara: Crystalizing the q -analogue of universal enveloping algebras, *Comm. Math. Phys.*, **133**, (1990), 249-260.
- [22] M. Kashiwara: On crystal bases of the q -analogue of universal enveloping algebras, *Duke Math. J.*, **63**, (1991), 465-516.
- [23] K.C. Misra and T. Miwa: Crystal base for the basic representation of $U_q(\hat{sl}(n))$, *Comm. Math. Phys.*, **134**, (1990), 79-88.
- [24] M. Jimbo, K.C. Misra, T. Miwa and M. Okado: Combinatorics of representations of $U_q(\hat{sl}(n))$ at $q = 0$, *Comm. Math. Phys.*, **136**, (1991), 543-566.
- [25] O. Foda and T. Miwa: Corner transfer matrices and quantum affine algebras, *Int. J. Mod. Phys. A*, **7**, Suppl. 1A, (1992), 279-302.
- [26] B. Davies, O. Foda, M. Jimbo, T. Miwa and A. Nakayashiki: Diagonalization of the XXZ Hamiltonian by vertex operators, *Comm. Math. Phys.*, **151**, (1993), 89-153.

- [27] M. Idzumi, K. Iohara, M. Jimbo, T. Miwa, T. Nakayashima and T. Tokihiro: Affine symmetry in vertex models, *Int. J. Mod. Phys. A*, **8**, (1993), 1479-1511.
- [28] E. Date and M. Okado: Calculation of excitation spectra of the spin model related with the vector representation of the quantized affine algebra of type $A_n^{(1)}$, *Osaka Univ. Math. Sci. Preprint*, **1**, (1993).
- [29] M. Jimbo, T. Miwa and Y. Ohta: Structure of the space of states in RSOS models, *Int. J. Mod. Phys. A*, **8**, (1993), 1457-1477.
- [30] M. Jimbo, K. Miki, T. Miwa and A. Nakayashiki: Correlation Functions of the XXZ model for $\Delta < -1$, *Phys. Lett. A*, **168**, (1992), 256-263.
- [31] O. Foda, M. Jimbo, K. Miki, T. Miwa and A. Nakayashiki: Vertex operators in solvable lattice models, *J. Math. Phys.*, to appear.
- [32] M. Jimbo: Introduction to the Yang-Baxter equation, *Int. J. Mod. Phys. A*, **4**, (1989), 3759-3777.
- [33] Y. Koyama: Staggered Polarization of Vertex models with $U_p(\mathfrak{sl}(n))$ Symmetry, *Comm. Math. Phys.*, (1993), in press.
- [34] I. B. Frenkel and N. Yu. Reshetikhin: Quantum affine algebras and holonomic difference equations, *Comm. Math. Phys.*, **149**, (1992), 1-60.
- [35] P. Goddard, A. Kent and D. Olive: Virasoro algebra and coset space models, *Phys. Lett. B*, **152**, (1985), 88-92.
 — : Unitary representations of the Virasoro and super-Virasoro algebras, *Comm. Math. Phys.*, **103**, (1986), 105-119.
- [36] R. J. Baxter: Solvable eight-vertex model on an arbitrary planar lattice, *Phil. Trans. Roy. Soc. Lond.*, **289**, (1978), 315-346.
- [37] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press (1982).
- [38] M. Jimbo, *Nankai Lecture Notes on Mathematical Physics*, ed. Ge Mo-Lin, World Scientific (1992).
- [39] B. Davies: Corner transfer matrices and quantum affine algebras, *J. Phys. A: Math. Gen.*, (1993), in press.
- [40] B. Davies: Corner transfer matrices for the Ising model, *Physica A*, **154**, (1989), 1-20.
- [41] B. Davies and P. A. Pearce: Conformal invariance and critical spectrum of corner transfer matrices, *J. Phys. A: Math. Gen.*, **23**, (1990) 1295-1312.
- [42] K. Sogo and M. Wadati: Boost operator and its application to quantum Gelfand-Levitan equation for Heisenberg-Ising chain with spin one-half, *Prog. Theor. Phys.*, **69**, (1983), 431-450.
- [43] H. B. Thacker: Corner transfer matrices and Lorentz invariance on a lattice, *Physica D*, **18**, (1986), 348-359.
- [44] R. J. Baxter: Spontaneous staggered polarization of the F model, *J. Stat. Phys.*, **9**, (1973), 145-182.

- [45] E. Date, M. Jimbo, K. Miki and M. Okado: Mean staggered polarization for the higher spin analog of the 6-vertex model, *Int. J. Mod. Phys. A*, **7**, *Suppl.* 1A, (1992), 151-160.
- [46] M. Jimbo, T. Miwa and A. Nakayashiki: Difference equations for the correlation functions of the eight-vertex model, *J. Phys. A: Math. Gen.*, **26**, (1993), 2199-2209.
- [47] R. J. Baxter and S. B. Kelland: Spontaneous polarization of the eight vertex model, *J. Phys. C: Solid State Phys.*, **7**, (1974), L403-406.